

## **Soils Map**





State: lowa

County: **Montgomery** Location: **3-71N-37W** 

Township: Scott
Acres: 60.2
Date: 9/24/2020





	ymbol: IA137, S						,				,					
Code	Soil Description	Acres	Percent of field	CSR2 Legend	Non- Irr Class *c	*i Corn	*i Alfalfa	*i Soybeans	*i Bluegrass	*i Tall Grasses	CSR2**	CSR	*n NCCPI Overall	*n NCCPI Corn	*n NCCPI Small Grains	*n NCCPI Soybeans
9D2	Marshall silty clay loam, 9 to 14 percent slopes, eroded	18.03	30.0%		Ille	177.6	5	51.5	3.2	5.3	61	58	86	86	63	6
822D2	Lamoni clay loam, 9 to 14 percent slopes, eroded	12.79	21.2%		IVe	100.8	2.6	29.2	1.8	3	7	15	59	59	54	4
9C2	Marshall silty clay loam, 5 to 9 percent slopes, eroded	11.11	18.5%		Ille	211.2	5.9	61.2	3.8	6.3	87	68	91	91	67	7
Z24E2	Shelby clay loam, deep loess, 14 to 18 percent slopes, eroded	6.29	10.4%		IVe	0	0	0	0	0	38		71	71	51	5
Z24D2	Shelby clay loam, deep loess, 9 to 14 percent slopes, eroded	5.46	9.1%		IIIe	0	0	0	0	0	52		76	76	55	51
9B	Marshall silty clay loam, 2 to 5 percent slopes	3.81	6.3%		lle	232	6.5	67.3	4.2	7	95	85	89	89	74	7



11B	Ackmore- Colo-Judson	2.71	4.5%		llw	200	4.2	58	3.6	6	79	68	86	84	52	82
	complex, 2 to 5 percent															
	slopes															
	Weighted Average				erage	137.3	3.7	39.8	2.5	4.1	54.1	*-	*n 78.9	*n 78.8	*n 60	*n 60.6

<sup>\*\*</sup>IA has updated the CSR values for each county to CSR2.

\*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.

\*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.